

Provinces Panama.

Habitat: Found in both lowland and montane forests in Eastern Panama. Highly arboreal, spending most of their lives in an around epiphytes.

Size: Relatively small frog 17-20mm svl



Veraguas +Coclé **Province**



























Red headed harlequin poison frog

Oophaga anchicayensis "Red head

Distribution: Valle de Cauca Colombia in the region of San José de Anchicaya.

Habitat: Common in Primary and secondary rainforests as well as some altered habitats, almost always below 750m asl.

Size: Relatively small morph 30-35mm svl



www.fantasticfrogs.co.ul

Anchicaya harlequin poison frog

Oophaga anchicayensis "Anchicaye"

Distribution: Valle de Cauca Colombia in the region of San José de Anchicaya.

Habitat: Common in Primary and secondary rainforests as well as some altered habitats, almost always below 750m asl.

Size: large morph 35—45mm svl



Valle De Cauca Colombia

www.fantasticfrogs.co.ul

Pacasi harlequin poison frog

Oophaga anchicayensis "Pacasi"

Distribution: believed to be found in the region of San José de Anchicaya.

Habitat: Common in Primary and secondary rainforests as well as some altered habitats, almost always below 750m asl.

Size: large morph 35—45mm svl





Fuego harlequin poison frog

Oophaga sp. "Fuego"

Distribution: Actual origin unknown

Habitat: Common in Primary and secondary rainforests as well as some altered habitats, almost always below 750m asl.

Size: large morph 35-45mm svl





Bullseye harlequin poison frog

Oophaga histrionica "Red and Black Bullseye"

Distribution: Area surrounding Santa

Cecilia, department of Riseralda, Colombia.

Habitat: Found in a narrow geographic range at an altitude of 340-410m asl in Primary rainforest.

Size: Average sized morph 30-40mm svl

Department de Riseralda, Colombia

































Oophaga histrionica "Blue"

Distribution: Area surrounding Santa Cecilia, department of Riseralda, Colombia.

Habitat: Found in a narrow geographic range at an altitude of 340-410m asl in **Primary rainforest.**

Size: Average sized morph 30-40mm svl

Department de Riseralda, Colombia





























Atrato harlequin poison frog

Oophaga histrionica "Atrato"

Distribution: Area surrounding Santa

Cecilia, department of Riseralda, Colombia.

Habitat: Found in a narrow geographic

range at an altitude of 340-410m asl in

Primary rainforest.

Size: Average sized morph 30-40mm svl

Department de Riseralda, Colombia

































Pacurita harlequin poison frog

Oophaga andresi "Quibido"

Distribution: Area surrounding Pacurita east of the Atrato river, Colombia.

Habitat: Found in lowland rainforests along the eastern bank of the Atrato river from 55–168m temperatures are consistently warm

Size: Average sized morph 33—43mm svl



Valle de Cauca Colombia



Pacurita harlequin poison frog

Oophaga andresi "Yellow brown blotched"

Distribution: Area surrounding Pacurita east of the Atrato river, Colombia.

Habitat: Found in lowland rainforests along the eastern bank of the Atrato river from 55–168m temperatures are consistently warm


Size: Average sized morph 33—43mm svl













Lehmans harlequin poison frog

Distribution: Found near Dagua west of the Anchicaya river, Colombia.

Habitat: Found in lowland rainforests along the Anchicaya river district temperatures are consistently warm

VULNERABLE ENDANGERED

Size: Average sized morph 33-43mm svl







Lehmans harlequin poison frog

Distribution: Found near Dagua west of the Anchicaya river, Colombia.

Habitat: Found in lowland rainforests along the Anchicaya river district temperatures are consistently warm

VULNERABLE ENDANGERED

Size: Average sized morph 33—43mm svl





Diabolito poison frog

Oophaga sylvatica"Paru

Distribution: Found from San Lorenzo east towards Lita, Ecuador.

Habitat: Found in lowland rainforests as well as some altered habitats where the humidity remains high year round.



Ecuado www.fantasticfrogs.co.uk

/ Diabolito poison frog

Oophaga sylvatica "Bilsa"

Distribution: Found from Puerto Quito South towards Santa Domingo, Ecuador.

Habitat: Found in lowland rainforests as well as some altered habitats where the humidity remains high year round.

Size: Average sized morph 26—38mm svl





Diabolito poison frog

Oophaga sylvatica "San Lorenzo"

Distribution: Found close to San Iorenzo, Ecuador.

Habitat: Found in lowland rainforests as well as some altered habitats where the humidity remains high year round.

VULNERABLE ENDANGERED

Size: Average sized morph 26—38mm svl

LEAST





Diabolito poison frog

Oophaga sylvatica"Lita"

VULNERABLE ENDANGERED

CRITICALLY EXTINCT ENDANGERED IN THE WILD

Distribution: Found from Lita Ecuador west towards Alto Tambo.

Habitat: Found in lowland rainforests as well as some altered habitats where the humidity remains high year round.

Size: Average sized morph 26—38mm svl

NEAR THREATENED

LEAST

